

OG05 Hysteroscopic Endometrial Resection

Expires end of February 2023

You can get information locally from:

Huddersfield Royal Infirmary

- Main switchboard, who can connect you to the relevant department, on 01484 342 000
- Patient Advice and Liaison Service (PALS) on 01484 342 128

Calderdale Royal Hospital

- Main switchboard, who can connect you to the relevant department, on 01422 357 171
- Patient Advice and Liaison Service (PALS) on 01422 222 417

You can also contact:

You can get more information from www.aboutmyhealth.org

Tell us how useful you found this document at www.patientfeedback.org



Royal College
of Surgeons
of England



THE ROYAL
COLLEGE OF
SURGEONS
OF EDINBURGH



Patient Information Forum

eidohealthcare.com

EIDO
HEALTHCARE

UNITED KINGDOM

What is an endometrial resection?

An endometrial resection is an operation to remove the lining (endometrium) of your uterus (womb). After the operation most women have a noticeable reduction in their periods and, for some women, periods stop altogether.

Your gynaecologist has suggested an endometrial resection. However, it is your decision to go ahead with the operation or not. This document will give you information about the benefits and risks to help you to make an informed decision.

If you have any questions that this document does not answer, it is important that you ask your gynaecologist or the healthcare team. Once all your questions have been answered and you feel ready to go ahead with the procedure, you will be asked to sign the informed consent form. This is the final step in the decision-making process. However, you can still change your mind at any point.

What are the benefits of surgery?

The most common reason for having an endometrial resection is to relieve the symptoms of heavy periods (abnormal uterine bleeding). An endometrial resection is another effective treatment instead of a hysterectomy. It also has fewer complications and a quicker recovery time.

An endometrial resection may be suitable if your heavy periods are caused by one or more of the following conditions.

- Fibroids, where the muscle of your womb becomes overgrown.
- Polyps – A polyp is an overgrowth of the lining of your womb that looks like a small grape on a stalk.
- Excessive thickening of the lining of your womb.

For most women, no specific cause can be found for heavy periods.

Most women will have much less bleeding when they have their period. Pain is usually significantly reduced, although for some women mild cramping may still happen.

About a third of women who have the operation will not have periods anymore.

Are there any alternatives to surgery?

Heavy periods can be treated using a variety of non-hormonal and hormonal oral (by mouth) medications.

Other alternatives include an IUS (intra-uterine system - an implant containing a synthetic form of the hormone progesterone that fits in your womb) but these are usually tried before surgery is recommended.

You should discuss the options with your gynaecologist.

What will happen if I decide not to have the operation?

Your doctor will continue to try to control your symptoms with medication, or you can continue without treatment.

For some women this is acceptable if the cause of the symptoms is not serious.

What happens before the operation?

Depending on your age and symptoms, your gynaecologist may recommend that you have a biopsy (removing small pieces of tissue from the lining of your womb). Your gynaecologist may also recommend an ultrasound scan or hysteroscopy (a procedure to look at the inside of your womb using a small telescope).

Your gynaecologist may ask you to take a hormonal treatment for up to 3 months before the operation. The treatment will make it easier to remove the lining of your womb by thinning it and reducing the size of any polyps or fibroids. You may be able to have the operation straight after a period to avoid the need for hormonal treatment.

Your gynaecologist may arrange for you to have a pre-admission assessment. They will carry out several tests and checks to find out if you are fit enough for the operation. If you have any questions about the operation, you should ask the healthcare team at this visit.

Your gynaecologist may ask you to have a pregnancy test. Sometimes the test does not show an early-stage pregnancy so let your gynaecologist know if you could be pregnant.

What does the operation involve?

The healthcare team will carry out a number of checks to make sure you have the operation you came in for. You can help by confirming to your gynaecologist and the healthcare team your name and the operation you are having.


The operation is usually performed under a general anaesthetic but various anaesthetic techniques are possible. Your anaesthetist will discuss the options with you. You may also have injections of local anaesthetic to help with the pain after the operation.

You may be given antibiotics during the operation to reduce the risk of infection.

The operation usually takes about 30 minutes.

Your gynaecologist will examine your vagina. They will pass a small telescope (operating hysteroscope or resectoscope) through your vagina, across your cervix (neck of your womb) into your womb.

Your gynaecologist will pass fluid through the telescope to distend (swell) your womb. They will use a diathermy resecting loop (a loop of wire heated by electricity) to remove the lining of your womb and any polyps or small fibroids they find.



A hysteroscopy

What should I do about my medication?

Make sure your healthcare team knows about all the medication you take and follow their advice. This includes all blood-thinning medication as

well as herbal and complementary remedies, dietary supplements, and medication you can buy over the counter.

How can I prepare myself for the operation?

If you smoke, stopping smoking now may reduce your risk of developing complications and will improve your long-term health.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight.

Regular exercise should help to prepare you for the operation, help you to recover and improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.

If you have not had the coronavirus (Covid-19) vaccine, you may be at an increased risk of serious illness related to Covid-19 while you recover. Speak to your doctor or healthcare team if you would like to have the vaccine.

What complications can happen?

The healthcare team will try to reduce the risk of complications.

Any numbers which relate to risk are from studies of people who have had this operation. Your doctor may be able to tell you if the risk of a complication is higher or lower for you. Some risks are higher if you are older, obese, you are a smoker or have other health problems. These health problems include diabetes, heart disease or lung disease.

Some complications can be serious and can even cause death.

You should ask your doctor if there is anything you do not understand.

Your anaesthetist will be able to discuss with you the possible complications of having an anaesthetic.

General complications of any operation

- Feeling or being sick. Most women have only mild symptoms and feel better within 1 to 2 days without needing any medication.

- Bleeding or discharge, lasting up to 4 weeks. It starts off heavy but gradually gets lighter.
- Infection (risk: 3 in 100). Most infections are minor and often happen after leaving hospital. They are usually easily treated with antibiotics.
- Allergic reaction to the equipment, materials or medication. The healthcare team is trained to detect and treat any reactions that might happen. Let your gynaecologist know if you have any allergies or if you have reacted to any medication or tests in the past.
- Blood clot in your leg (deep-vein thrombosis – DVT) (risk: less than 1 in 200). This can cause pain, swelling or redness in your leg, or the veins near the surface of your leg to appear larger than normal. The healthcare team will assess your risk. They will encourage you to get out of bed soon after the operation and may give you injections, medication, or special stockings to wear. Let the healthcare team know straight away if you think you might have a DVT.
- Blood clot in your lung (pulmonary embolus), if a blood clot moves through your bloodstream to your lungs. Let the healthcare team know straight away if you become short of breath, feel pain in your chest or upper back, or if you cough up blood. If you are at home, call an ambulance or go immediately to your nearest Emergency department.
- Chest infection. If you have the operation within 6 weeks of catching Covid-19, your risk of a chest infection is increased (see the 'Covid-19' section for more information).

Specific early complications

- A small hole in your womb made by one of the instruments, with possible damage to a nearby structure (risk: 1 in 100). You may need to stay overnight for close observation in case you develop complications. If your gynaecologist is concerned that an organ has been damaged, you may need keyhole surgery or another operation involving a larger cut.
- Bleeding during the operation (risk: 1 in 100). The risk is higher if you have fibroids. The bleeding can prevent the procedure from

being completed or lead to you needing a hysterectomy (risk: 1 in 500).

- Fluid overload, where the fluid used to expand your womb enters your bloodstream (risk: less than 2 in 100). You may need to stay overnight for close observation and you may be given medication.
- Failed procedure, if it is not possible to place the instruments into your womb.

Specific late complications

- Haematometra, where blood and other menstrual fluid collect in pockets in your womb (risk: less than 1 in 100). If these pockets do not drain across your cervix or through your fallopian tubes, they can cause pain. Most women will not have periods and the pockets are usually noticed on a scan.
- Blood and fluid collecting in your fallopian tubes and becoming trapped (risk: less than 1 in 1,000). This causes pain.
- Continued bleeding or pain needing further treatment. This may involve a hysterectomy (risk: 3 in 10 in the first 10 years).

Covid-19

A recent Covid-19 infection increases your risk of lung complications or death if you have an operation under general anaesthetic. This risk reduces the longer it is since the infection. After 7 weeks the risk is no higher than someone who has not had Covid-19. However, if you still have symptoms the risk remains high. The risk also depends on your age, overall health and the type of surgery you are having.

You must follow instructions to self-isolate and take a Covid-19 test before your operation. If you have had Covid-19 up to 7 weeks before the operation you should discuss the risks and benefits of delaying it with your surgeon.

Consequences of this procedure

- Pain is a cramping pain similar to a period and is usually easily controlled with simple painkillers such as paracetamol.

How soon will I recover?

In hospital

After the operation you will be transferred to the recovery area and then to the ward.

The healthcare team will tell you what was found during the operation and discuss with you any treatment or follow-up you need.

You should be able to go home the same day. However, your doctor may recommend that you stay a little longer.

If you are worried about anything, in hospital or at home, contact the healthcare team. They should be able to reassure you or identify and treat any complications.

Returning to normal activities

If you had sedation or a general anaesthetic and you do go home the same day:

- a responsible adult should take you home in a car or taxi and stay with you for at least 24 hours;
- you should be near a telephone in case of an emergency;
- do not drive, operate machinery or do any potentially dangerous activities (this includes cooking) for at least 24 hours and not until you have fully recovered feeling, movement and co-ordination; and
- do not sign legal documents or drink alcohol for at least 24 hours.

To reduce the risk of a blood clot, make sure you carefully follow the instructions of the healthcare team if you have been given medication or need to wear special stockings.

You may get some cramps and mild bleeding similar to a period. Rest for 1 to 2 days and take painkillers if you need them.

You should be able to return to normal activities after 2 to 4 days. Most women are fit for work after about a week.

You should expect to have some bleeding or discharge for up to 4 weeks. This may be heavy and red to start with but will change to a red-brown discharge.

Use sanitary pads, not tampons. To reduce the risk of infection, do not have sex, or have a bath or swim until the discharge has settled.

Let your doctor know if you develop any of the following problems.

- A high temperature.
- Heavy bleeding or an unpleasant-smelling discharge from your vagina.
- Your pain does not settle or increases and is not relieved by your medication.
- Pain in your lower leg.
- Breathing difficulties.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Do not drive until you can control your vehicle, including in an emergency, and always check your insurance policy and with the healthcare team.

Will I need hormone replacement therapy (HRT)?

An endometrial resection will not affect when you go through menopause. At the time of menopause, if you want to go on HRT, your doctor should give you an oestrogen and progesterone hormone.

Do I still need smear tests?

As the operation has no effect on your cervix, continue to have regular smear tests.

Will I still be able to have children?

The operation is not recommended for women who still want children. Serious complications for you and your baby can happen if you become pregnant after an endometrial resection.

Even if your periods stop, there is still a risk of becoming pregnant. Do not rely on the operation as a method of contraception. If neither you nor your partner has been sterilised, continue to use a reliable method of contraception.

Summary

An endometrial resection is a common gynaecological operation. It helps relieve the symptoms of heavy periods. You should get less bleeding and pain.

Surgery is usually safe and effective but complications can happen. You need to know about them to help you to make an informed decision about surgery. Knowing about them will also help to detect and treat any problems early.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you, and they may be able to tell you about any other suitable treatments options.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

Acknowledgements

Reviewer

Jeremy Hawe (MBChB, MRCOG)

Illustrator

Medical Illustration Copyright © Nucleus Medical Art. All rights reserved. www.nucleusinc.com